



# ISOLATION AND IDENTIFICATION OF MARINE PHYTOPLANKTONS FOR PRODUCTION OF CARBOHYDRATE TYPE BIOMASS

Syahrudin Kasim <sup>[a]</sup> and M. Sjahrul <sup>[a]\*</sup>

**Keywords:** Types of sea phytoplankton, Biomass production, Carbohydrate content, Phytoplankton culture, Arschat medium

The primary aim of the present study was to isolate and identify a number of sea phytoplanktons for biomass production and determination of carbohydrate content. Sea phytoplankton was cultivated in cultured media "Ares-chat" to multiply seeds and cultivated stocks. In the mass media cultivation, isolation and identification of sea plankton was done based on size and some specific characteristics. The measurement of temperature, salinity and pH was also done in the media. The biomass weight of the density of selected phytoplankton was done by gravimetric method and the carbohydrate content was determined by glucose with Luff School method. The isolation results indicate six different types of sea phytoplanktons: *Chlorella sp.*, *Dunaliella sp.*, *Tetraselmis chuii*, *Chaetoceros calcitrans*, *Chaetoceros gracilis*, and *Chaetoceros Isocrysis galbana*. The highest biomass content 0.34 gL<sup>-1</sup> was on *Chlorella sp.* and the lowest 0.14 gL<sup>-1</sup> was on *Isocrysis galbana*. The carbohydrate content was also varied, the highest 31.99% on *Dunaliella sp.* and the lowest 4.70% on *Chaetoceros gracilis*.

\* Corresponding Authors

E-Mail: eddievedder\_jie@yahoo.com

[a] Department of Chemistry, University of Hasanuddin F.MIPA, Makassar, Indonesia


## Introduction

Microsize water organisms floating with the water movement are known as plankton.<sup>1</sup> Plankton concentration at the surface of sea water varies from 500 cells/mL to 10 cells mL.<sup>2</sup> Plankton plays an important role and sometimes dominates material cycle in sea.<sup>3</sup> Planktons are divided into two groups: phytoplankton and zooplankton.<sup>4,5</sup>

Most phytoplankton are very small to see by the naked eye, but large in number. They appear as green in color in water due to chlorophyll in their cells, although the real color is varied for each species due to the presence of an extra pigment such as phycobiliprotein.<sup>6,7</sup> Phytoplankton usually gathers at the euphotic zone where light intensity that makes photosynthesis possible (Arinardi et al., 1997 and Richtel, M., 2007). Phytoplankton gets energy by photosynthesis,<sup>8,9</sup> for which they must be on the sea surface (called as euphotic zone), lake or other source of water. Through photosynthesis, phytoplankton produces a lot of oxygen into the earth's atmosphere.<sup>10</sup> Because of the photosynthesis, the organism lives on the surface of water, and are also able to produce carbohydrate (glucose).<sup>11,12</sup> The ability of phytoplankton to synthesise its organic material, make them to be the main source of food chain in the sea ecosystem and fresh water (Fig.1)<sup>12</sup>

This planktonic organism is usually collected by using a net. Based on the size of mesh of the net, phytoplankton is classified as megaplankton (bigger than 0.2 mm;) and macro plankton (0.2-2.0 mm).

Microplankton is sizes 20-0.2 mm.<sup>13</sup> A plankton which is captured by filter milipore is known nanoplankton, whereas



a very small plankton sizes 2-20  $\mu\text{m}$ , ultraplankton, a smaller plankton less than 2  $\mu\text{m}$ .<sup>14</sup>

Indonesia has the longest coastal line in the world which is about  $\pm$  80.791.42 km. Phytoplankton consists of 30.000 species.<sup>15</sup> Its habitat is on the surface of different types of water bodies. It is a large perspectives of phytoplankton utilization for production of carbohydrate as biofuel (bioethanol and biobutanol) and raw material to substitute the food grade corn or cereal raw materials.<sup>16</sup> The other benefit of phytoplankton is that they are able to absorb carbon dioxide and converse it to oxygen. As much as 90% of dry weight of phytoplankton adsorbe carbon dioxide so that it is able to reduce gas up to 1.000 tons/ha/year.<sup>17</sup>

**Table 1.** Macronutrients (A) and micronutrients (B) for marine alga production

Macro Nutrients								
No.	Mineral Salt	c, 10 <sup>-3</sup> mol L <sup>-1</sup>	g L <sup>-1</sup>	Note	Added			
					g	mL L <sup>-1</sup> )		
1.	NaCl	300	25.38	direct addition				
2.	MgSO <sub>4</sub> .7H <sub>2</sub> O	10	4.730	direct addition				
3.	KNO <sub>3</sub>	4	0.503	stock 1 M/100 mL	10.1	3		
4.	KH <sub>2</sub> PO <sub>4</sub>	2	0.236	stock 1 M/100 mL	13.6	1		
5.	CaCl <sub>2</sub> .6H <sub>2</sub> O	15	1.210	stock 1 M/1000 mL	110.99	10		
Micronutrients								
No.	Mineral salt	c, 10 <sup>-6</sup> mol.L <sup>-1</sup>	Stock 1 M mL <sup>-1</sup>	g	Added mL L <sup>-1</sup>	Stock 2 M mL <sup>-1</sup>	Added, mL Stock1 Stock 2 <sup>+</sup> *	
6.	H <sub>3</sub> BO <sub>3</sub>	300	0.4/100	2.5724	1	-		
7.	ZnSO <sub>4</sub> .7H <sub>2</sub> O	1	0.2/100	3.7274		0.02/100	1	1
8.	MnSO <sub>4</sub> .4H <sub>2</sub> O	1	0.4/100					
9.	CuSO <sub>4</sub> .5H <sub>2</sub> O	0.2	0.01/100	0.3495		0.01/100	1	1
10.	CoCl <sub>2</sub> .6H <sub>2</sub> O	0.2	0.01/100	0.3738		0.01/100	1	1
11.	(NH <sub>4</sub> ) <sub>2</sub> MoO <sub>4</sub> .4H <sub>2</sub> O	0.2	0.01/100	0.8016		0.01/100	1	1
12.	NaFeEDTA	25						
13.	Na <sub>2</sub> SiO <sub>3</sub> .9H <sub>2</sub> O**	15	0.1/100	2.6909	1	-		
14.	Tris mix (7.8): TRIS-HCl TRIS-base	0.023% (w/w); mean: 0.025 g Tris mix in 100 g water (MJ water = 1 g.mL <sup>-1</sup> )			0.01722 0.00778			
15.	Vitamin mix: Biotin Vitamin B1 Vitamin B12				1 0.2 0.15 0.2			

Notes: \* Use 1 mL of stock solution to make a stock solution of 2; + Use 1 mL of stock solution 2 to make 1 L medium; \*\* Use for diatom

## Experimentals

### Materials and methods

Seeds extracted from pure cultures Pitoplankton Research Institute of Fisheries and Marine Maros, South Sulawesi.

### Instrument and Apparatus

Equipment cups: Consists of the culture vessel, covering 1000 mL bottle size, tank size of 60 liters and a set of glassware. Salinity was measured salinity meter.

### Identification of types of sea plankton with high carbohydrate content

#### Isolation and identification of phytoplanktons

Isolation of phytoplankton in the present study was done by taking samples of mass culture result for shrimp feed which contains bacteria and/or other microorganism such as Protozoa. In order to get pure phytoplankton strain, washing the cell at the new culture media and planted in medium to be selected for study. The culture medium used was modification of several culture media with artificial sea water which is known as "Ars-chat" which stands for Arifin-

Syaharul-Syahruddin-Henk Schat. Whereas the jelly medium was made by mixing the gelatin in medium solution of Ars-chat as much as 1% (10 grams). After autoclaved, it was poured into a gelatin plate each about 20 ml. When the gelatin plate is cool, 3 drops are spread over the surface of gelatin plate by a sterile pipette. After a span of 3 to 5 days the growth of phytoplankton cell colony was observed and found that phytoplankton did not associate with colony of bacteria. Further, 2-5 cells of phytoplankton were implanted in a test tube containing Ars-chat for phytoplankton culture of about 5-10 mL. After a span of 8 – 15 days the phytoplankton cell would grow with high density (10<sup>3</sup> – 10<sup>6</sup> cells/mL). Then the seed of this phytoplankton cell was planted in the medium Ars-chat of which the volume was more (in Erlenmeyer glass of 50 mL, 250 mL, and 500 mL) then in culture bottle 1000 mL.

Further culture was done to multiply the seed aside from being kept for culture stock.

#### Physico-chemical parameters of water samples

In order to find out the growth condition of phytoplankton to be isolated, measurement of several physico-chemical parameters was done, such as temperature, salinity, and pH in situ in mass culture (in the field).




Figure 2. Research method

### Pure Culture of Phytoplankton

Pure culture of phytoplankton was initiated by inoculating seed of phytoplankton cell into a culture tube with density of low phytoplankton cell in Erlenmeyer flask of 500 mL. After observing for 8 days, the culture condition was seen to be old enough (density ranges from 105-106 sel/mL) the culture was divided into 3 parts. The first and second parts each 200 mL was put into a culture bottle volume 1 liter. Whereas the rest 100 mL was added with medium Ars-chat as much as 400 mL, to make it to 500 mL culture volume in erlenmeyer flask as culture stock again kept for 8 days in the future. Whereas the 1 Liter volume culture after 8 days was shifted to aquarium volume of 60 liters. For both culture bottles of 1 liter, each of its growth pattern was observed daily for ten days by using Haemositometer through microscope. At this culture the flow of its specific growth was observed using the formula:

$$\mu = \frac{\ln N_t - \ln N_0}{t} \quad (1)$$

where

- $N_t$  is the density of the cell population at the time  $t$  (cells mL<sup>-1</sup>)  
 $N_0$  is the density of the cell population at baseline (cells mL<sup>-1</sup>)  
 $\mu$  is the specific growth rate constant (hr<sup>-1</sup>)  
 $t$  is the time (hours)

The amount of biomass in each liter culture volume was observed by filtering culture medium during the optimum growth by using a suitable filter paper as per the size of phytoplankton. The weight of phytoplankton was set by weighing the filter paper used before and after.

### Biomass culture of selected phytoplankton types and Its weight determination.

Referring to physico-chemical parameter, the growth condition of phytoplankton in field measured in situ, selected phytoplankton was cultured in large amount at aquarium of 60 liters. After the density of optimal phytoplankton, 1 liter was taken out for the mass culture to determine its biomass gravimetrically. Then harvesting process was done by stopping the light and aeration for about 3 days for the separation between water and phytoplankton by sedimentation.

### Analyzing carbohydrate contents of selected phytoplankton.

Carbohydrate analysis was done by using glucose reduction determination method of Luff-Schoorl.

## Results and Discussion

### Isolation and Identification of sea phytoplankton



This study has been carried out for the isolation and identification of a number of marine phytoplanktons from both, dinoflagellate and diatom groups: Dinoflagellate consists of *Chlorella sp.*, *Dunaliella sp.*, and *Tetraselmis chuii*, diatom species of *Chaetoceros calcitrans* consist of, *Chaetoceros gracilis* and *Chaetoceros Isocrysis Galbana* (Table 2).

Observation result on the growth of flagellate types: *Chlorella sp.*, *Dunaliella sp.*, and *Tetraselmis chuii*, in artificial water medium (Fig.3). Observation result of growth pattern of diatom *Chaetoceros calcitrans* and *chaetoceros gracilis* in artificial water medium (Fig. 4).

The growth pattern of the three phytoplanktons viz. *Chlorella sp.*, *Dunaliella sp.* and *Tetraselmis chuii* cultured at experimental environmental conditions, such as culture temperature, continuous light, nutrient used, salinity of sea water for each species of phytoplankton. Water pH applied and aeration CO<sub>2</sub>, show growth pattern similar to one which underwent 3 stages of growth, namely adjustment, splitting and growth-dead. Optimum density was attained on the seventh day for phytoplankton type *Chlorella sp.* and for *Tetraselmis chuii* it was attained on the sixth day as 1.238.333 x 10<sup>4</sup> cell mL<sup>-1</sup> and 173.250 x 10<sup>4</sup> cell mL<sup>-1</sup>. (Table 4), respectively.

**Table 2.** Results of isolation and identification of phytoplanktons

No.	Types of Phytoplankton	Size ( $\mu\text{m}$ )	Information
1.	<i>Chlorella sp.</i>	3-4	Green, not motile and no flagella. Spherical cell with cup-shaped chloroplasts
2.	<i>Dunaliella</i>	Width 5-8 Length 7-12	Green motile with two flagella, near the back of the cells, The cells move quickly in the water with shaking behaviour while swimming. Cells are spherical to cylindrical and usually have a red-eye point.
3.	<i>Tetraselmis chuii</i>	width 9-10 Length 12-15	Motile green, with a four flagella grew out of a groove in the back of the anterior cells. The cells move rapidly in the water and looked shaken while swimming. There are four lobes are elongated and have a point reddish eyes.
4.	<i>Isochrysis galbana</i>	4 – 8	Motile cells with two flagella that grow near the rear part of the cell. Cells move faster in the water and turning while swimming. These algae are spherical shaped, gold-colored and usually have a red-eye point.
5.	<i>Chaetoceros gracilis</i>	0,5–2	These organisms are single cells and may form an interconnected chain using a hook of the adjacent cell. The main body is like a petri dish shape. Side view of square-shaped organisms gives an impression of spines from the corner.
6.	<i>Chaetoceros calcitrans</i>	4	Cylindrical and reddish brown in colour

**Figure 3.** Growth pattern of dinoflagellate species *Chlorella sp.*, *Dunaliella sp.*, and *Tetraselmis chuii*.**Figure 4:** The pattern of growth of Diatoms *Chaetoceros calcitrans* and *Chaetoceros gracilis*.

Specific growth rate ( $\mu$ ) of phytoplankton types *Chlorella sp.*, *Dunaliella sp.*, and *Tetraselmis chuii* (Table 5).

The pattern of specific growth rate ( $\mu$ ) of diatom group for types *Chaetoceros calcitrans* and *Chaetoceros gracilis* cultured at experimental environmental conditions as stated earlier in the present paper. Optimum density of both phytoplanktons was attained on the sixth day for phytoplankton types *Chaetoceros calcitrans*,  $713.333 \times 10^4$  cells per mL and  $223.667 \times 10^4$  cell per mL respectively (Table 6).

**Table 4.** Phytoplankton density of *Chlorella sp.*, *Dunaliella sp.*, *Tetraselmis chuii*, at the 8<sup>th</sup> day of culture

Time (days)	Types of Phytoplankton density, cells mL <sup>-1</sup>		
	<i>Chlorella sp.</i> (10 <sup>5</sup> )	<i>Dunaliella sp.</i> (10 <sup>4</sup> )	<i>T. chuii</i> (10 <sup>4</sup> )
0	10,000	25,000	10.000
1	19,583	69,333	13.167
2	92,000	412,667	32.333
3	257,000	844,333	80.333
4	453,833	1.126,500	130.667
5	627,333	1.204,667	161.333
6	654,000	1.238,333	173.250
7	656,333	1.158,667	150.667
8	634,000	1.089,333	131.255

**Table 6.** Density and *Chaetoceros calcitrans* phytoplankton *Chaetoceros gracilis* cultures on 8<sup>th</sup> day

Time, days	Types of phytoplankton density, cells mL <sup>-1</sup>	
	<i>Chaetoceros calcitrans</i> (10 <sup>4</sup> )	<i>Chaetoceros gracilis</i> (10 <sup>4</sup> )
0	10,000	2,500
1	42,167	10,333
2	206,667	51,667
3	331,667	112,000
4	533,333	174,667
5	660,000	198,333
6	713,333	223,667
7	650,000	216,667
8	538,333	214,333

Specific growth rate ( $\mu$ ) of phytoplankton types *Chaetoceros calcitrans* and *Chaetoceros gracilis* can be seen in Table 7.

**Table 5.** Phytoplankton cell density and specific growth rate ( $\mu$ )

Species	Size $\mu\text{m}$	t, days	Observations, cells mL <sup>-1</sup>		$\mu, \text{d}^{-1}$
			N <sub>0</sub>	N <sub>t</sub>	
<i>Chlorella sp.</i>	3–4	6	1.000.000	66.000.000	0.0318
<i>Dunaliella sp.</i>	8	6	250.000	13.500.000	0.0302
<i>Tetraselmis chuii</i>	10–15	7	100.000	1.725.000	0.0198

Table 7. Phytoplankton cell density and specific growth rate ( $\mu$ )

Species	Size ( $\mu\text{m}$ )	t, days	Observation cells mL <sup>-1</sup>		$\mu, \text{d}^{-1}$
			N <sub>0a</sub>	N <sub>t</sub>	
<i>Chaetoceros calcitrans</i>	4	6	100.000	7.133.333	0.0291
<i>Chaetoceros gracilis</i>	0,5–2	6	25.000	2.236.667	0.441

#### Biomass culture and determination of dry weight of selected phytoplankton

Determination of the relationship between biomass dry weight of phytoplankton and culture volume in optimum phase (Table 8).

**Table 8.** Relationships of phytoplankton biomass dry weight and 1 liter of culture volume in the optimum phase

No.	Types of phytoplankton	Biomass dry weight, mg L <sup>-1</sup>
1.	<i>Chlorella sp.</i>	0,34
2.	<i>Dunaliella sp.</i>	0,28
3.	<i>Tetraselmis chuii</i>	0,33
4.	<i>Chaetoceros calcitrans</i>	0,27
5.	<i>Chaetoceros gracilis</i>	0,24
6.	<i>C. Isocrysis galbana</i>	0,14

To predict the biomass obtained from mass culture of phytoplankton, dry weight test was done. Based on dry weight test of biomass of phytoplankton cultured with Arschat medium, it was observed that for phytoplankton of dinoflagellata type was having higher dry weight as compared to phytoplankton of diatom type. From the data it can be concluded that to obtain biomass in large amount, it is suggested to culture the type of dinoflagellata with dry weight as much as 0.34 mg/L, 0.33mg/L, and 0.28 mg/L mass culture medium respectively for phytoplanktons *Chlorella sp.*, *Tetraselmis chuii*, and *Dunaliella sp.*

#### Carbohydrate analysis of selected phytoplankton

The results of carbohydrate test contained in each sea phytoplankton are recorded in Table 9.

**Table 9.** Carbohydrate levels on various types of species of phytoplankton biomass in the culture.

No.	Types of phytoplankton	Carbohydrate levels (%)
1.	<i>Chlorella sp.</i>	30,75
2.	<i>Dunaliella sp.</i>	31,99
3.	<i>Tetraselmis chuii</i>	26,68
4.	<i>Chaetoceros calcitrans</i>	6,02
5.	<i>Chaetoceros gracilis</i>	4,70
6.	<i>C. Isocrysis galbana</i>	12,00

On the basis of the type of cultured phytoplankton, it is apparent to note that phytoplankton which has higher percentage of carbohydrate content is from dinoflagellata group, and that are of phytoplankton types *Dunaliella sp.*, *Chlorella sp.* and *Tetraselmis chuii*, which have carbohydrate content up to 31.99%, 30.75% and 26.68% respectively.

## Conclusions

Isolation results for six types of phytoplanktons are obtained. They are *Chlorella sp.*, *Dunaliella sp.*, *Tetraselmis chuii*, *Chaetoceros calcitrans*, *Chaetoceros gracilis* and *Chaetoceros Isocrysis galbana*. The highest biomass 0.34 g/L is of *Chlorella sp.* of dinoflagellata type and the lowest is 0.14 g/L of *Isocrysis galbana* of diatom type, whereas the highest carbohydrate content is in *Dunaliella sp.* i.e. 31.99% of dinoflagellata type and the lowest is in *Chaetoceros gracilis* i.e. 4.70% of diatom type. Phytoplankton with high carbohydrate content is suggested to be processed further to become vegetable fuel.

## References

- <sup>1</sup>Arinardi, O.H., A.B. Sutomo, S.A. Joseph, Trianingsih, E. Asnaryanti and S. H. Riyono.. *Plankton Abundance composition range and predominant in Indonesia East Region at Territorial of the Sea*. P3O-LIPI. Jakarta. **1997**
- <sup>2</sup>Bu-Olayan, A.H., Al-Hassan, R., Thomas, B.V., Subrahmanyam, M. N., *Environ. Int.* **2001**, 26(4), 199.
- <sup>3</sup>Brand Larry E., Sunda William G., and Guillard R. R. L.. *J.Exp. Marine Biol. Ecol.*, **1986**, 96(3), 225.  
[http://dx.doi.org/10.1016/0022-0981\(86\)90205-4](http://dx.doi.org/10.1016/0022-0981(86)90205-4)
- <sup>4</sup>Bold, H. C., Wynne, M. J. *Introduction to the algae: Structure and Reproduction*, 2nd Ed., Prentice-Hall. Inc., New Jersey **1985**.
- <sup>5</sup>Botes. L, *Phytoplankton Identification Catalogue*, Globallast Monograph Series No.7. IMO London, Saldanha Bay, South Africa, **2001**.
- <sup>6</sup>Cermeno, P., de Vargas, C., Abrantes, F., Falkowski, P. G. *PLOS ONE* **2010**, 5(4), e10037.
- <sup>7</sup>Jobman, D., *Will ethanol produce a bull market in Corn?* Chicago Board of Trade, **2006**.  
<http://www.cbot.com/cbot/docs/74305.pdf>
- <sup>8</sup>Fenchel. T., *Ann. Rev. Ecol. Syst.*, **1988**, 19.
- <sup>9</sup>Joint, I., Jordan, M. B., *J. Plymouth Marine Lab.*, **2008**, 30(2), 199.
- <sup>10</sup>*Methods of Biochemical Analysis and Food Analysis using Test Combination*, Boehringer Mannheim GmbH Biochemica, Germany, **1987**.
- <sup>11</sup>Miljanovic, B., Mijic, I., Jovanovic, V., Sipos, S., *Response of Phytoplankton Association of Zavojsko Reservoir to Seasonal Variation in Hydrochemical Parameters*, Balwois, Ohrid, Republic of Macedonia, **2010**, No.25.
- <sup>12</sup>Morel, F. M. M., *J. Compilation Geobiol.*, **2008**, 6, 318.
- <sup>13</sup>Newell, G. E., Newell, R. C., *Marine plankton: A practical guide*, Hutchinson Educational LTD Great Portland Street, London, 1963, vol.1. 178-202.
- <sup>14</sup>Parker D. L., Rai, L. C., Mallick, N., Rai, P. K., Kumar, H. D., *J. Appl. Environ. Microbiol.*, **1988**, 64(4), 1545.
- <sup>15</sup>Richtel. M, *The New York Times*, 01 May, **2007**
- <sup>16</sup>Thurman. H. V, *Introductory Oceanography*, Prentice Hall College, New Jersey, USA, 1997; Kotai, L., Gacs, I., *Eur. Chem. Bull.*, **2012**, 1(1), 1;
- <sup>17</sup>Yamaji. I, *Illustrations of the marine plankton of Japan*. Hoikusha publishing Co., Ltd. **1982**.

Received: 15. Sept. 2012.

Accepted: 09. Oct. 2012

